

KINGDOM OF SAUDI ARABIA MINISTRY OF NATIONAL GUARD - HEALTH AFFAIRS KING ABDULAZIZ MEDICAL CITY - RIYADH DEPARTMENT OF ONCOLOGY


# Hickman Line

### Tips before inserting a central catheter:

- Mak sure to meet the patient educator to explain the procedure and to answer any related questions before the insertion of the catheter.
- The catheter (Hickman Line) is part of your treatment plan, thus it is very important to take care of it in order to reduce and prevent any subsequent complications.
- You must take a shower and wipe your body with antiseptic wipes (please consult with your nurse) before the procedure.
- Inform your treating physician if you have any signs and symptoms of sickness before the procedure.
- You can continue your work and daily activities. Make sure that you take a shower daily. Avoid heavy activities and prevent any trauma to the catheter site or any accidental pulling of the catheter.


### What is Hickman Line?

- A Hickman Line is a flexible rubbery central venous catheter that is attached to one of the neck's major vessels.
- The central venous catheter is inserted to relieve frequent tingling as well as some treatments that can only be administered through the main blood veins through the venous catheter.
- The line will be inserted in the Interventional Radiology Department under local anesthesia. It will be removed in the same department, in the patient's room or in outpatient clinics at the end of the treatment plan.



### **Aim of Hickman Line:**

- Administration of chemotherapy.
- Administration of medications.
- Transfusion of blood and blood components.
- Blood draw.



### Care for Hickman line area:

- Wash your hands before touching the catheter or the surrounding area.
- Maintain the catheter dressing, personal hygiene and shower bearing in mind the catheter should be covered with water proof dressing.
- Make sure to change the tie, which holds the catheter lumens when needed and after shower.
- After 48 hours of insertion visit the clinic to change the dressing when sterile gauze dressing is applied, and after 7 days when a transparent dressing is used.
- Inspect the catheter exit site for signs of bacterial infection like redness, swelling, and discharge. Visit the Emergency Department in the main hospital (KAMC) immediately if any sign appeared.
- Always Keep the catheter clamps closed when not in use.
- Maintain the catheter by having the dressing changed and the catheter lumen flushed on a weekly basis in the clinic or according to the physician instructions.
- Know when to call for help and when to go to the Emergency Department.

### **Precautions while Hickman line in:**

- Protect the catheter, so it will not be pulled out of its position accidentally.
- Swimming in pools and in bathtubs is not allowed.
- In case the dressing is loose, proceed to the clinic during weekdays from 8:00 AM to 4:00 PM or to the Emergency Department in the main hospital (KAMC) after 4 PM, prefereble to call help line phone number before coming to the clinic (0118040909) .



## Emergencies that might occur with Hickman line at home and how to manage them

Emergency:	How to manage:
- Sudden pain in the neck, chest or shoulders. - A cough or shortness of breath.	<ul> <li>- Make sure the lumens are locked.</li> <li>Lay down on the opposite side of the catheter.</li> <li>- If the pain persists, proceed to the Emergency Department in the main hospital (KAMC).</li> </ul>
- Accidental removal of the catheter.	- Press on the exit site for five minutes with sterile gauze (it will be provided by the nursing team before discharge) and proceed to the Emergency Department in the main hospital (KAMC).
- Swelling in face, neck, chest or arms.	- Proceed to the Emergency Department in the main hospital (KAMC).
- Discharge, redness, swelling, or bleeding from the catheter exit site.	- Proceed to the Emergency Department in the main hospital (KAMC).
- Rising body temperature or chills.	- Measure your temperature. If it is 38 °C or more proceed to the Emergency Department in the main hospital (KAMC).  - Do not use any antipyretics.  - If the temperature is less than 38°C, continue to measure it every hour.
- Any crack or opening in the catheter lumens or any leakage.	- Close the lumens lock between the opening site and your chest then proceed to the Emergency Department in the main hospital (KAMC).

# Bring your yellow card with you to the Emergency Department in the main hospital (KAMC) and show it to the staff there.



### هـذا المريـض يستـخدم عـلاج كيـميائي



#### Take your Alert card and go to The ER immediately if:

- ✓ Fever of °38C or higher (don't take any fever lowering medication).
- ✓ Shivering or general unwell feeling.
- ✓ Bleeding that doesn't stop anywhere in the body or unusual bruises or skin rash.
- ✓ Continues vomiting & not being able to eat or drink or continues diarrhea.
- ✓ Redness or oozing around the central line.
- ✓ Increasing of mouth sores severity especially if prevented you from eating and drinking.

#### حضر هذه البطاقة وتوجه للطوارئ في حال:

- ✓ الاصابة بحرارة ٣٨ درجة مئوبة فما فوق (لا تتناول مخفض للحرارة).
  - ✓ الشعور برجفة أو تعب جسدى شديد.
  - ✓ ألم في الصدر وصعوبة في التنفس.
  - ✓ نزيف لا يتوقف من أي مكان في الجسم أو كدمات غير مألوفة أو طفح جلدى.
  - ✓ قيء مستمر وعدم القدرة على الأكل أو الشرب أو إسهال مستمر.
    - ✓ إحمرار أو قيح حول القسطرة المركزية.
    - ✓ ازدباد تقرحات الفم خاصة اذا منعتك من الاكل والشرب.



رقم خدمه المساعدة والتواصل لمرضى الأورام وذويهم

Tel: 0118040909

هاتف: ۹۰۹۰۹۰۱۰

يكن الإتصال من الساعة (٨) صباحاً إلى( ٥ )مساءً فقط أثناء أوقات الدوام الرسمي

### Don't hesitate to call help line:

- •If you have any questions about your treatment.
- •If you have any other queries, patient and family education.

Please bring your medications during doctor visit

## www.ngha.med.sa

2022